

SYSTAFLAM

Diclofenac Diethylamine, Mephenesin, Menthol, Methyl Salicylate, Turpentine Oil

COMPOSITION:

Systaflam Gel: Each gram gel contains Diclofenac Diethylamine BP 0.0116 gm, Mephenesin IP 0.05 gm, Menthol BP 0.02 gm, Methyl Salicylate BP 0.05 gm and Turpentine Oil BP 0.03 gm.

PHARMACOLOGY:

Systaflam Gel (Diclofenac Diethylamine, Mephenesin, Menthol, Methyl Salicylate, Turpentine oil) works by reducing pain and delayed onset muscle soreness; relieving pruritus; providing relief from painful conditions; providing relief of musculoskeletal aches and pains; reducing the pain.

INDICATION:

Systaflam Gel is a Non-steroidal anti-inflammatory (NSAID) drug used to treat pain associated with conditions like gout, migraine, rheumatoid arthritis, sprains of muscles and joints etc.

DOSAGE AND ADMINISTRATION:

Depending on the size of the area to be treated, 2-4 gm of **Systaflam Gel** should be applied to the skin 3-4 times daily to the affected area and rubbed in lightly.

CONTRAINDICATION:

It is contraindicated in patients who have hypersensitivity to **Systaflam Gel**. In addition, **Systaflam Gel** should not be used if one has the following conditions:

- children less than 12 years old
- cuts
- infected skin

SIDE EFFECTS:

The most commonly reported side-effects of **Systaflam Gel** are itching, blisters, allergic contact dermatitis, anaphylactic reactions, pruritus, and rash.

PRECAUTIONS:

Systaflam Gel should not be allowed to come in contact with the eyes or mucous membranes. After application, the hands should be washed unless they are the sites being treated. Not to be taken by mouth.

USE IN PREGNANCY AND LACTATION:

Not recommended.

PHARMACEUTICAL PRECAUTION:

It should be stored in a cool and dry place, protected from light and moisture.

HOW SUPPLIED:

Systaflam Gel: Each tube contains 15 gm/25 gm/30 gm of gel.

Manufactured by:



MEDICON Pharmaceuticals Ltd.
Mirpur, Dhaka, Bangladesh